

primary studies - published RCT

Nebulized gentamicin in children with cystic fibrosis: Enhancing antibiotic lung deposition by increasing flow rate and fill volume.

Code: CN-00195584 **Year:** 1997 **Date:** 2001

Author: Mallo J

Study design (if review, criteria of inclusion for studies)

Randomised, cross-over trial.

Participants

13 participants with CF with severe lung disease. Mean (SD) age 26 (5.9) years. Mean (SD) FEV1 % predicted, 31.7(10.6); awake PaO₂ range 53-77 mmHg; PaCO₂ 45 mmHg; mean (SD) BMI 20 (3) kgm².

Interventions

Order of intervention randomised. Night 1: Room air and low-level CPAP (4 - 5 cm H₂O). Night 2: Oxygen (1.4 +/- 0.9L/min to maintain SaO₂ 90%) and low-level CPAP (4 - 5 cm H₂O). Night 3: BVS +/- oxygen (0.7+/-0.9 L/min to maintain SaO₂ 90%). 3 nights within a 1-week period. Time between nights unclear.

Outcome measures

VI, VT; RR; respiratory disturbance indices; SaO₂ TcCO₂.

Main results

During RA and LFO₂ studies, patients wore a nasal mask with a baseline continuous positive airway pressure (CPAP) of 4 to 5 cm H₂O. Minute ventilation (V I) was measured using a pneumotachograph in the circuit and was not different between wake and non-rapid eye movement (NREM) sleep on any night. However, V I was reduced on the RA and LFO₂ nights from awake to rapid eye movement (REM) (p

Authors' conclusions

BVS leads to improvements in alveolar ventilation during sleep in this patient group.

<http://dx.doi.org/10.1089/jam.1997.10.331>

See also

Journal of Aerosol Medicine: Deposition, Clearance, and Effects in the Lung YR: 1997 VL: 10 DE: RCT NO: 4

Keywords

Adult; Artificial Ventilation; non pharmacological intervention - devices OR physiotherapy; Oxygen; Ventilators; Sleep Disorders;