

primary studies - published RCT

Evaluation of once daily tobramycin versus the traditional three time daily for the treatment of acute pulmonary exacerbations in adult cystic fibrosis patients.

Code: CN-00613134 **Year:** 2006 **Date:** 2010

Author: Al Ansari NA

Study design (if review, criteria of inclusion for studies)

randomized double-blind, placebo-controlled study

Participants

20 patients with CF. One individual failed to conclude the study. All subjects were clinically stable when studied and without recent infectious exacerbation or hospitalization in the last 30 days.

Interventions

Groups were randomized for placebo (n = 10; mean age 12.1 +/- 6.0) or 3 mg melatonin (n = 9; mean age 16.6 +/- 8.26) for 21 days.

Outcome measures

Actigraphy was performed for 6 days before the start of medication and in the third week (days 14-20) of treatment. Isoprostane and nitrite levels were determined in exhaled breath condensate (EBC) at baseline (day 0) and after treatment (day 21).

Main results

Melatonin improved sleep efficiency (P = 0.01) and tended to improve sleep latency (P = 0.08). Melatonin reduced EBC nitrite (P = 0.01) but not isoprostane. In summary, melatonin administration reduces nitrite levels in EBC and improves sleep measures in clinically stable CF patients.

Authors' conclusions

The failure of melatonin to reduce isoprostane levels may have been a result of the low dose of melatonin used as a treatment.

<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/134/CN-00613134/frame.html>

See also

Qatar Medical Journal YR: 2006 VL: 15 DE: RCT NO: 1

Keywords

Adolescent; Adult; Child; Hormones; Melatonin; Mental Disease-Psychiatric Conditions; Nitrites; pharmacological_intervention; placebo; Sleep Disorders;