

primary studies - published RCT

## Treatment with ivacaftor in CF patients with the G551D mutation is associated with improvement in cognition

Code: CN-01608969    Year: 2018    Date: 2010

Author: Wilson J

### Study design (if review, criteria of inclusion for studies)

RCT

### Participants

10 CF and 11 control subjects

### Interventions

Subjects inhaled an aerosol containing nonabsorbable technetium-99m sulfur colloid (Tc-SC) particles and In-DTPA.

### Outcome measures

absorptive clearance rate of DTPA

### Main results

Tc-SC clearance from the lung is exclusively mucociliary, while In-DTPA is cleared by both absorption and mucociliary clearance. The difference between the In-DTPA and Tc-SC clearance rates estimates In-DTPA absorption. Tc-SC (mucociliary) clearance was similar in central and peripheral zones in CF and non-CF lungs. Total In-DTPA clearance was increased in both zones in CF lungs. The absorptive component of In-DTPA clearance was increased in the airway-dominated central lung zones in CF ( $42\% \times h^{-1}$ ) versus  $32\% \times h^{-1}$ ,  $p = 0.03$ ). The absorption of In-DTPA is increased in the CF airway.

### Authors' conclusions

Further study is needed to understand the relative roles of fluid absorption, inflammation and other mechanisms potentially affecting epithelial permeability and DTPA absorption.

<https://www.cochranelibrary.com/central/doi/10.1002/central/CN-01608969/full>

### See also

Journal of cystic fibrosis

### Keywords

Adolescent; Adult; Biomarker; Inhalation OR nebulised;