

primary studies - published RCT

Effects of hyperoxia on oxygen uptake kinetics in cystic fibrosis patients as determined by pseudo-random binary sequence exercise.

Code: PM10029341

Year: 1999 **Date:** 2003

Author: Kusenbach G

Study design (if review, criteria of inclusion for studies)

randomized clinical trial

Participants

Patients (n = 39) who were at least 15 years of age and diagnosed with CF participated in the intervention

Interventions

3-session written self-disclosure intervention. Participants in the intervention arm were asked to write in the health care setting about an important emotionally distressing issue of personal significance for a 20-min period of time and two additional 20-min writing episodes at the patient's home, which were prompted by telephone calls. Patients in the control condition received standard care alone.

Outcome measures

number of days patients spent in the hospital, FEV1, BMI, subjective markers of health status.

Main results

Findings revealed that the intervention resulted in a reduction of the number of days patients spent in the hospital over a 3-month period. The intervention did not have an impact on physiological (Forced Expiratory Volume and Body Mass Index) or subjective markers of health status.

Authors' conclusions

These findings extend those of Pennebaker's (cf., J. Smyth, 1998) demonstrating an effect of the written-self-disclosure intervention on health care utilization. These preliminary findings are promising and justify further investigation of the modified intervention in other chronic illness populations.

<http://dx.doi.org/10.1007/s004210050494>

See also

Eur J Appl Physiol Occup Physiol. 1999 Jan;79(2):192-6.

Keywords

Adolescent; Adult; Hospitalization; Hospital care; non pharmacological intervention - psycho-soc-edu-org; Home; Organization;