

---

*primary studies - published RCT*

## **A prospective treatment trial of nasal polyps in adults with cystic fibrosis.**

**Code:** PM10953842

**Year:** 2000 **Date:** 2000

**Author:** Hadfield PJ

### **Study design (if review, criteria of inclusion for studies)**

Randomised, double blind, placebo-controlled. Parallel design.

### **Participants**

46 adults (over 16 years old) with CF and nasal polyps, excluding those who were pregnant or breast feeding, taking oral steroids, taking more than 1500 micrograms of inhaled steroid per day, had a severely deviated nasal septum or had undergone a surgical

### **Interventions**

22 participants received the active form of the drug with only 10 completing the course. 24 participants were prescribed the placebo and 11 of this group completed. Nasal drops in the form of either betamethasone sodium diphosphate nasal drops or passive placebo drops containing an identical vehicle, prescribed as two drops to be used twice a day for 6 weeks, in a head down and forwards position.

### **Outcome measures**

Primary outcome was polyp size. Secondary outcomes were subjective symptom scores.

### **Main results**

betamethasone nasal drops showed a statistically significant reduction in polyp size in comparison to placebo.

<http://www.rhinologyjournal.com/>

### **See also**

Rhinology. 2000 Jun;38(2):63-5.

### **Keywords**

Adult; Betamethasone; Intranasal; Nasal Polyps; pharmacological\_intervention; Respiratory Tract Diseases; Steroids; Anti-Inflammatory Agents;