

primary studies - published RCT

Low-flow oxygen and bilevel ventilatory support: effects on ventilation during sleep in cystic fibrosis.

Code: PM11208637

Year: 2001 Date: 2005

Author: Milross MA

Study design (if review, criteria of inclusion for studies)

Single centre randomised placebo controlled study.

Participants

18 CF patients (5.5-36.3 years, median 15.1). Diagnosis not clear, mean sweat chloride, 74.5 mg/L. One patient had 5T mutation (associated with a milder phenotype).

Interventions

Azithromycin (10 patients) 250 mg twice a week for 12 weeks versus placebo (8 patients).

Outcome measures

BPI-ANCA levels in the blood (a possible marker of inflammation). Also respiratory function. Also secondary outcomes not fully reported (authors contacted). Overall weight gain in both groups, but wide range and differences not reported.

Main results

BPI-ANCA was found in 12 CF patients (67%) and four (22%) healthy subjects (P

Authors' conclusions

BPI-ANCA levels are higher among patients colonised with *P. aeruginosa*. Twelve weeks of AZM therapy did not lower the BPI-ANCA level in patients with CF.

<http://ajrccm.atsjournals.org/content/163/1/129.full.pdf+html>

See also

Am J Respir Crit Care Med. 2001 Jan;163(1):129-34.

Keywords

Adolescent; Adult; Anti-Bacterial Agents; Azithromycin; Child; pharmacological_intervention; placebo; Proteins; Bacterial Infections; Respiratory Tract Infections; Respiratory Tract Diseases; Infection; Pseudomonas aeruginosa; Pseudomonas; Macrolides; Anti-Inflammatory Agents; Anti-Inflammatory Agents - excl Steroids;