

primary studies - published RCT

## Efficacy of once-daily tobramycin monotherapy for acute pulmonary exacerbations of cystic fibrosis: a preliminary study.

Code: PM11340683

Year: 2001 Date: 2001

Author: Master V

### Study design (if review, criteria of inclusion for studies)

double-blind RCT, 2 years.

### Participants

CF patients with PA-induced pulmonary exacerbations. Conv group (n = 51 admissions); Mono group (n = 47)

### Interventions

CF patients received either once-daily tobramycin (Mono) or conventional therapy with tobramycin/ceftazidime given 8-hourly (Conv). Tobramycin doses were adjusted to achieve a daily area under the time-concentration curve of 100 mg x hr/L in both groups.

### Outcome measures

Results were assessed for both short-term changes (efficacy and safety after 10 days of IV antibiotics during acute exacerbations) and long-term changes (efficacy, safety, and sputum microbiology between study entry and exit). Pulmonary function tests (PFTs) on admission were similar in both groups.

### Main results

After 10 days of IV antibiotics, absolute mean improvements in percent of predicted PFTs were 12.8, 12.1, and 13.7 for forced expiratory volume in 1 sec (FEV(1)), forced vital capacity (FVC), and forced expired flow between 25--75% of FVC (FEF(25--75%)) in the Conv group (n = 51 admissions) compared to 10.6, 9.9, and 10.6 in the Mono group (n = 47)(P<0.05 for all). Sixteen percent in the Conv group and 15% of patients in the Mono group did not respond to therapy by day 10. Long-term PFT patterns were similar for the Conv and Mono groups. The time between admissions did not differ. The Mono group showed a significant increase in tobramycin minimum inhibitory concentrations (MICs) against PA from study entry to study exit (P = 0.02, n = 27 strains); this failed to reach significance in the Conv group (P = 0.08, n = 25). There was no significant increase in the number of isolates, with MIC> or =8 mg/L in both groups. No short- or long-term changes in audiology or serum creatinine were found in either group. After 10 days of IV therapy, the urinary enzyme N-acetyl-beta-d-glucosaminidase/creatinine ratios increased in both groups (P0.05). This increase was greater in the Conv compared to the Mono group (P

### Authors' conclusions

this pilot study indicates once-daily tobramycin therapy to be as effective and safe as conventional 8-hourly tobramycin/ceftazidime therapy. Combination antibacterial therapy appears to offer no clinical advantage over once-daily tobramycin monotherapy. Tobramycin once-daily monotherapy is a potential alternative to conventional IV antibacterial therapy which deserves further investigation, including the impact on susceptibility of PA to tobramycin.

<http://dx.doi.org/10.1002/ppul.1060>

### See also

Pediatr Pulmonol. 2001 May;31(5):367-76.

### Keywords

Adolescent; Adult; Anti-Bacterial Agents; Bacterial Infections; Ceftazidime; Cephalosporins; Child; Combined Modality Therapy; Delayed-Action Preparations; Infection; Intravenous; Monotherapy; pharmacological\_intervention; Pseudomonas aeruginosa; Pseudomonas; Respiratory Tract Diseases; Respiratory Tract Infections; Tobramycin; Exacerbation; Aminoglycosides;