

primary studies - published RCT

## **Dose-finding and 24-h monitoring for efficacy and safety of aerosolized Nacystelyn in cystic fibrosis.**

**Code:** PM11866009

**Year:** 2002 **Date:** 2006

**Author:** App EM

### **Study design (if review, criteria of inclusion for studies)**

RCT

### **Participants**

15 patients (mean age 12.5 y, range 6.9-21.5 y) participated.

### **Interventions**

The treatments were randomized and performed on 2 separate occasions, 8 weeks apart.

### **Outcome measures**

Spirometry was conducted before and after each treatment. Transcutaneous oxygen tension (P(tO<sub>2</sub>)) was measured.

### **Main results**

There were no changes in spirometry values. During PEP, different trends in blood-gas tension were seen, and there were no consistent changes. During oscillating PEP, P(tO<sub>2</sub>) increased and P(tCO<sub>2</sub>) decreased. During oscillating PEP, P(tCO<sub>2</sub>) was lower and the intra-individual change in P(tCO<sub>2</sub>) was more pronounced than during PEP. The results obtained immediately after oscillating PEP showed a higher P(tO<sub>2</sub>) and a lower P(tCO<sub>2</sub>) than with PEP.

### **Authors' conclusions**

PEP and oscillating PEP can both cause transitory effects on blood gases in patients with cystic fibrosis. However, oscillating PEP alters blood-gas tensions more than does PEP, and hyperventilation during oscillating PEP may reduce treatment time.

<http://dx.doi.org/10.1183/09031936.02.00025802>

### **See also**

Eur Respir J. 2002 Feb;19(2):294-302.

### **Keywords**

Adolescent; Adult; Airway clearance technique; Chest Wall Oscillation; Child; non pharmacological intervention - devices OR physiotherapy; oscillating devices; Positive-Pressure Respiration- PEP- pep mask; Chest physiotherapy;