

primary studies - published RCT

Anti-inflammatory effects of montelukast in mild cystic fibrosis.

Code: PM12487226

Year: 2002 Date: 2005

Author: Schmitt-Grohé S

Study design (if review, criteria of inclusion for studies)

RCT

Participants

Participants were admitted to hospital for acute exacerbation. All participants performed HFCWO 1 - 3 times daily as outpatients before admission, but none had performed PEP. 15 participants, 8 males, 7 females. Aged at least 7 years, mean (SD) age 17.5 (4.2) years.

Interventions

Treatment lasted 30 minutes. PEP versus HFCWO. Both treatments were alternated within 48 hours of hospital admission and then reversed prior to discharge.

Outcome measures

RFTs and SaO₂ measured before and after every intervention. Each intervention was only done twice i.e. day 1 or 2 following admission then day -1 or -2 prior to discharge.

Main results

At admission and discharge, PEP breathing increased SpO₂ during treatment, whereas HFCWO decreased SpO₂ during treatment. Ventilation distribution, gas mixing, and lung function improved after HFCWO or PEP breathing.

Authors' conclusions

High-frequency chest wall oscillation and PEP breathing are similarly efficacious in improving ventilation distribution, gas mixing, and pulmonary function in hospitalized people with CF. Because SpO₂ decreases during HFCWO, people who have moderate to severe CF and who use HFCWO should have SpO₂ monitored during an acute exacerbation.

[http://dx.doi.org/10.1016/S1081-1206\(10\)62108-4](http://dx.doi.org/10.1016/S1081-1206(10)62108-4)

See also

Ann Allergy Asthma Immunol. 2002 Dec;89(6):599-605.

Keywords

non pharmacological intervention - devices OR physiotherapy; Airway clearance technique; Exacerbation; Respiratory Tract Infections; Respiratory Tract Diseases; Infection; Bacterial Infections; Positive-Pressure Respiration- PEP- pep mask; High Frequency Chest Wall Oscillation -HFCWO-; VEST Airway Clearance System; oscillating devices; Chest physiotherapy;