
primary studies - published RCT

Immunogenicity of a new purified fusion protein vaccine to respiratory syncytial virus: a multi-center trial in children with cystic fibrosis.

Code: PM12744878

Year: 2003 **Date:** 2003

Author: Piedra PA

Study design (if review, criteria of inclusion for studies)

phase II, multi-center, adjuvant-controlled trial

Participants

respiratory syncytial virus (RSV) seropositive children with cystic fibrosis (CF)

Interventions

vaccine or adjuvant-control

Outcome measures

RSV-specific, serum antibodies

Main results

At enrollment, RSV-specific, serum antibodies were comparable between both groups. At post-vaccination and end-of-study, RSV-specific, neutralizing antibody (Nt Ab) and binding antibody (Bd Ab) to the fusion (F) protein were significantly higher in PFP-3 vaccinees. After 28 days post-vaccination, Nt Ab and Bd Ab to F protein titers declined slowly at rates of 0.23 and 0.37 log₂ per month, respectively

Authors' conclusions

The PFP-3 vaccine-induced a robust immune response that lasted throughout the RSV season.

[http://dx.doi.org/10.1016/S0264-410X\(03\)00098-7](http://dx.doi.org/10.1016/S0264-410X(03)00098-7)

See also

Vaccine. 2003 Jun 2;21(19-20):2448-60.

Keywords

Child; fusion; Immunization; Infant; Infection; pharmacological_intervention; Proteins; Recombinant Proteins; Respiratory Syncytial Virus Infections; Respiratory Tract Diseases; Respiratory Tract Infections; Virus; Bronchiolitis;