

primary studies - published RCT

Bone mineral density in Australian children, adolescents and adults with cystic fibrosis: a controlled cross sectional study.

Code: PM14760157

Year: 2004 **Date:** 2008

Author: Buntain HM

Study design (if review, criteria of inclusion for studies)

RCT

Participants

16 CF patients terminally ill were randomized to telemedicine or standard care alone

Interventions

Patients in the telemedicine group had an ISDN line installed in their home and were given a videoconferencing unit connected to their home television set. Telemedicine patients had weekly videoconferences from home for a clinical assessment, psychological support and the opportunity for discussion with any member of the multidisciplinary team.

Outcome measures

Anxiety levels were measured before and after the conferences.

Main results

Eleven subjects completed the baseline assessment and seven patients completed the study (4 on telemedicine and 3 in the control arm). A total of 71 home videoconferences were conducted during the study. After six months there were no significant differences in quality of life, anxiety levels, depression levels, admissions to hospital or clinic attendances, general practitioner calls or intravenous antibiotic use between the two groups. However, there was a significant improvement in perception of body image for those in the telemedicine group and the patients liked and valued the service.

Authors' conclusions

The use of telemedicine can enhance the support that a specialist unit can provide for the patient and their family, and may reduce outpatient clinic attendances

<http://dx.doi.org/10.1136/thorax.2003.006726>

See also

Thorax YR: 2004 VL: 59 NO: 2

Keywords

Adult; Caregivers; Home; Home Care Services; Lung Transplantation; non pharmacological intervention - psycho-soc-edu-org; non pharmacological intervention - surg; telemedicine; transplantation; Depression; Mental Disease-Psychiatric Conditions; Organization;