

primary studies - published RCT

High-dose ibuprofen in cystic fibrosis: Canadian safety and effectiveness trial.

Code: PM17719932 **Year:** 2007 **Date:** 2007

Author: Lands LC

Study design (if review, criteria of inclusion for studies)

Multicenter double-blind placebo-controlled trial. Allocated treatment by a centralized pharmacy using a pre-defined block randomization schedule.

Participants

142 children with CF aged 6-18 years. Inclusion criteria: FEV1 >60% predicted at time of entry into the trial, with no hospitalizations in the previous 2 months. Exclusion criteria: people who had taken systemic corticosteroids or non-steroidal anti-inflammatory agents for more than 1 month in the past year, had abnormal hepatic, renal, hematologic disorders or coagulopathy, documented evidence of peptic ulcer di 18 participants (9 in each group) did not complete full 2 years of follow up, 11 due to adverse events (4 in treatment group, 7 in placebo group).

Interventions

All participants underwent a baseline pharmocokinetic study (baseline every hour for 3 hours), employing 200 mg tablets (Upjohn-Pharmacia) at a dose of 20 to 30 mg/kg to a maximum of 1600 mg.

Outcome measures

Annual rate of change in FEV1 % predicted, FVC % predicted, anthropometric data, chest radiograph score, number of hospitalizations (and length of stay), adverse effects, compliance, concomitant therapy (antibiotics, inhaled anti-inflammatory agents).

Main results

The patients in the high-dose ibuprofen group exhibited a significant reduction in the rate of decline of forced vital capacity percent predicted (0.07 +/- 0.51 vs -1.62 +/- 0.52; P = .03), but not FEV1%. The ibuprofen group also spent fewer days in hospital after adjusting for age (1.8 vs 4.1 days per year; P = .07). A total of 11 patients (4 in the ibuprofen group and 7 in the placebo group) withdrew due to adverse events.

Authors' conclusions

High-dose ibuprofen has a significant effect on slowing the progression of lung disease in CF and generally is well tolerated.

<http://dx.doi.org/10.1016/j.jpeds.2007.04.009>

See also

J Pediatr. 2007 Sep;151(3):249-54. Epub 2007 Jun 26.

Keywords

Adolescent; Anti-Inflammatory Agents; Child; High-Dose; Ibuprofen; pharmacological_intervention; Tablets; Anti-Inflammatory Agents - excl Steroids;