

---

primary studies - published RCT

## Safety, efficacy and convenience of tobramycin inhalation powder in cystic fibrosis patients: The EAGER trial.

**Code:** PM21075062

**Year:** 2011 **Date:** 2011

**Author:** Konstan MW

### Study design (if review, criteria of inclusion for studies)

open-label study, randomized

### Participants

553 patients aged  $\geq 6$  years

### Interventions

patients were randomized 3:2 to TIP (total 112mg tobramycin) via the Novartis T-326 Inhaler or TIS 300mg/5mL via PARI LC® PLUS nebulizer twice daily for three treatment cycles (28 days on-drug, 28 days off-drug).

### Outcome measures

Safety, efficacy, and treatment satisfaction outcomes were evaluated.

### Main results

TIP was generally well-tolerated; adverse events were similar in both groups. The rate of cough suspected to be study drug related was higher in TIP-treated patients (TIP: 25.3%; TIS: 4.3%), as was the overall discontinuation rate (TIP: 26.9%; TIS: 18.2%). Increases in FEV(1)% predicted from baseline to Day 28 of Cycle 3 were similar between groups; the mean reduction in sputum *P. aeruginosa* density (log(10) CFU/g) on Day 28 of Cycle 3 was also comparable between groups. Administration time was significantly less for TIP (mean: 5.6 versus 19.7min, p

### Authors' conclusions

TIP has a safety and efficacy profile comparable with TIS, and offers a far more convenient treatment option for pseudomonas lung infection in CF.

<http://dx.doi.org/10.1016/j.jcf.2010.10.003>

### See also

J Cyst Fibros. 2011 Jan;10(1):54-61. Epub 2010 Nov 12.

### Keywords

Adolescent; Adult; Aged; Anti-Bacterial Agents; Bacterial Infections; Child; Drug Administration Schedule; Infection; Inhalation OR nebulised; pharmacological\_intervention; Powders; Pseudomonas aeruginosa; Pseudomonas; Respiratory Tract Diseases; Respiratory Tract Infections; Tobramycin; Aminoglycosides;