

---

primary studies - published RCT

## Open randomised prospective comparative multi-centre intervention study of patients with cystic fibrosis and early diagnosed diabetes mellitus.

**Code:** PM24620855

**Year:** 2014 **Date:** 2014

**Author:** Ballmann M

### Study design (if review, criteria of inclusion for studies)

Multi-national, multicentre, open labelled, randomized and prospective controlled parallel group's trial

### Participants

The trial should include 74 subjects showing cystic fibrosis related diabetes newly diagnosed by oral glucose tolerance test during annual screening for cystic fibrosis related diabetes.

### Interventions

Repaglinide vs insulin injections.

### Outcome measures

Primary endpoint is mean HbA1c after 24 months of treatment. Secondary endpoints are change in FEV1% predicted and change in BMI-Z-score.

### Main results

The results regarding BMI after 6 months and 12 months showed an improvement for the insulin treated patients and were inconsistent for those treated with repaglinide. HbA1c and lung function (FEV1%pred) were unchanged for either group. The authors compared the changes -12 months to baseline and baseline to +12 months separately for each group. Therefore a direct comparison of the effect of repaglinide versus insulin on BMI, HbA1c and FEV1%pred was not presented.

### Authors' conclusions

There is only one prospective study comparing oral antidiabetic drugs to insulin in the treatment of CFRD without fasting hyperglycaemia.

<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3975280/>

### See also

BMC Pediatr. 2014 Mar 11;14:70. doi: 10.1186/1471-2431-14-70.

### Keywords

Adult; Aged; Child; Diabetes Mellitus; Gastrointestinal Diseases; Hypoglycemic Agents; Insulin; Pancreatic Diseases; pharmacological\_intervention;