

primary studies - published RCT

Pancreatic enzyme supplementation improves the incretin hormone response and attenuates postprandial glycaemia in adolescents with cystic fibrosis: a randomized crossover trial.

Code: PM24670086

Year: 2014 **Date:** 2017

Author: Perano SJ

Study design (if review, criteria of inclusion for studies)

Randomized, double blinded, cross-over pilot clinical trial.

Participants

13 patients with cystic fibrosis (CF) aged 6-23 years, 8 females,

Interventions

Subjects with CF received, in random order, oral glycine (0.5 g/kg/day, dissolved in any liquid) and placebo (glass sugar), each during 8 weeks with an intermediate 2-week wash-out period.

Outcome measures

Symptom scores. Spirometric variables. Pulse oximetry . TNF-alpha in serum and IL-6

Main results

As compared with placebo, after glycine intake patients had better symptom questionnaire scores ($p = 0.02$), mainly regarding sputum features and dyspnea. While spirometric variables tended to decline during placebo intake, they remained stable or even increased during glycine treatment ($p = 0.04$ to $p = 0.003$). In this context, FEV1 declined 8.6% after placebo and increased 9.7% at the end of the glycine period. Pulse oximetry improved after glycine intake ($p = 0.04$ vs. placebo). TNF-alpha in serum and IL-6 and G-CSF in sputum tended to decline at the end of the glycine period ($p = 0.061$, $p = 0.068$ and $p = 0.04$, respectively, vs placebo). Glycine was remarkably well tolerated.

Authors' conclusions

The clinical, spirometric and inflammatory status of subjects with CF improved after just 8 weeks of glycine intake, suggesting that this amino acid might constitute a novel therapeutic tool for these patients. Thus, further studies are warranted.

<http://dx.doi.org/10.1210/jc.2013-4417>

See also

J Clin Endocrinol Metab. 2014 Mar 26;jc20134417.

Keywords

Child; Glycine; non pharmacological intervention - diet; Supplementation; Amino Acids; Proteins;