

primary studies - published RCT

## **A randomized controlled trial of vitamin D replacement strategies in pediatric CF patients.**

**Code:** PM26211605

**Year:** 2016 **Date:** 2020

**Author:** Simoneau T

### **Study design (if review, criteria of inclusion for studies)**

Prospective cross-over trial

### **Participants**

Fourteen males and eight females from the national center for CF, with an average FEV(1) of 54%

### **Interventions**

Autogenic drainage (AD). Mechanical insufflator-exsufflator (MI-E). Subjects received either AD or MI-E in a random order.

### **Outcome measures**

Sputum was collected and weighed immediately after treatment. Subjects performed lung function tests at baseline, 20 minutes after and 1 hour after treatment; additionally, a 2-minute walk test was performed at the end of all lung function tests. Saturation, dyspnea scores while resting and after a 2-minute walk and subjective fatigue were recorded.

### **Main results**

Thirty-six percent more sputum was collected following MI-E than AD treatment (P<0.05).

### **Authors' conclusions**

Treatment with the MI-E was more effective for clearing sputum in CF subjects, initial evidence suggests that the MI-E may be successfully incorporated into treatment protocols. Further studies are needed to assess the long-term benefits of MI-E in patients with CF.

<http://dx.doi.org/10.1016/j.jcf.2015.07.004>

### **See also**

J Cyst Fibros. 2016 Mar;15(2):234-41. doi: 10.1016/j.jcf.2015.07.004. Epub 2015 Jul 23.

### **Keywords**

Adolescent; Airway clearance technique; Child; Drainage; non pharmacological intervention - devices OR physiotherapy; pharmacological\_intervention; Airway clearance drugs -expectorants- mucolytic- mucociliary-; Exacerbation; Respiratory Tract Infections; Respiratory Tract Diseases; Infection; Bacterial Infections; Postural Drainage; percussion; Chest physiotherapy; Autogenic Drainage;