
primary studies - published, non RCT

Safety and efficacy of prolonged levofloxacin inhalation solution (APT-1026) treatment for cystic fibrosis and chronic *Pseudomonas aeruginosa* airway infection.

Code: PM26935334

Year: 2016 **Date:** 2016

Author: Elborn JS

Study design (if review, criteria of inclusion for studies)

Open-label extension of a multinational, randomized study

Participants

Patients completing a multinational, randomized study comparing LIS and tobramycin inhalation solution (TIS)

Interventions

Levofloxacin inhalation solution (LIS) and tobramycin inhalation solution (TIS): three additional cycles of 28days of LIS 240mg twice daily followed by 28days off drug.

Outcome measures

Mean relative change in percent predicted forced expiratory volume in 1s (FEV1), time to pulmonary exacerbation, and patient-reported quality of life.

Main results

Extended treatment with LIS in 88 patients was well tolerated with no new safety signals and evidence of positive effects on FEV1 and quality of life.

Authors' conclusions

Patients receiving extended LIS treatment continued to show favorable efficacy with no additional safety concerns.

<http://dx.doi.org/10.1016/j.jcf.2016.01.005>

See also

J Cyst Fibros. 2016 Sep;15(5):634-40. doi: 10.1016/j.jcf.2016.01.005. Epub 2016 Feb 28.

Keywords

Adult; aeroquin; Aged; Anti-Bacterial Agents; Child; Inhalation OR nebulised; levofloxacin; pharmacological_intervention; Tobramycin; Bacterial Infections; Respiratory Tract Infections; Respiratory Tract Diseases; Infection; Quinolones; Aminoglycosides;