

primary studies - published RCT

Parent Experience With False-Positive Newborn Screening Results for Cystic Fibrosis.

Code: PM27485696

Year: 2016 **Date:** 1983

Author: Hayeems RZ

Study design (if review, criteria of inclusion for studies)

double-blind RCT

Participants

100 CF patients selected on the basis of clinical and bacteriological criteria divided were divided into five treatment groups

Interventions

Daily dosage of drugs, given i.v. as three divided doses, was as follows: penicillins 400 mg/kg, cephalosporins 200 mg/kg. Duration of therapy was 10-15 days.

Outcome measures

The side-effects were evaluated. The pharmacokinetics of three antibiotics (axiocillin, cefotaxime and cefoperazone) were also evaluated by determination of serum and sputum concentrations after single-dose injection during treatment. The antibiotics were tested in vitro by an agar dilution technique; MICs were assessed against strains of *Pseudomonas* spp. and *Pseudomonas aeruginosa* isolation from CF patients. Results were evaluated by means of clinical score, chest X-ray score and bacteriological score.

Main results

On the basis of in vitro testing, the highest sensitivity, both of *Pseudomonas aeruginosa* and *Pseudomonas* spp., was observed with ceftazidime, followed by piperacillin, azlocillin, cefsulodin and cefoperazone; cefotaxime and moxalactam showed lower activity rates and were therefore removed at an early stage from the trial. Preliminary clinical results indicated the following tendencies: 1) the best results were obtained with ceftazidime followed by cefsulodin and piperacillin; 2) *Pseudomonas* bacterial counts dropped considerably half way through the treatment and more at the end with ceftazidime and cefsulodin; with the other antibiotics there was a drop followed by a new rise at the end; 3) there was no close correlation among clinical, radiological and bacterial scores; 4) few side-effects were observed: a rise in temperature on the 10th-12th day with piperacillin, transient vomiting with cefsulodin and diarrhoea with cefoperazone. Moderate sputum levels of antibiotics were observed in all sputum fractions with ceftazidime but no antibiotic was detected with axlocillin; variable results were obtained with cefoperazone.

<http://dx.doi.org/10.1542/peds.2016-1052>

See also

Pediatrics. 2016 Aug 2. pii: e20161052.

Keywords

Anti-Bacterial Agents; Azlocillin; Bacterial Infections; Cefoperazone; Cefotaxime; Cefsulodin; Ceftazidime; Infection; Moxalactam; pharmacological_intervention; Piperacillin; *Pseudomonas aeruginosa*; *Pseudomonas*; Respiratory Tract Diseases; Respiratory Tract Infections; Penicillins; Cephalosporins;