

primary studies - published RCT

The role of small intestinal bacterial overgrowth in cystic fibrosis: a randomized case-controlled clinical trial with rifaximin.

Code: PM30232597

Year: 2018 **Date:** 2018

Author: Furnari M

Study design (if review, criteria of inclusion for studies)

Randomized controlled trial

Participants

79 CF patients (median age 19.6 years; 9.2-36.9) with a positive Glucose Breath Test (GBT)

Interventions

Rifaximin 1200 mg for 14 days or no treatment.

Outcome measures

Questionnaire and GBT were repeated 1 month after the end of treatment or 45 days after the first negative GBT. BMI, SDS-BMI, serum albumin levels (p

Main results

Out of 79 patients, 25 were affected by SIBO (31.6%) with a significant correlation with lower BMI, SDS-BMI ($p < 0.05$) and serum albumin levels ($p < 0.05$), independently from pancreas insufficiency. Twenty-three patients took part in the randomized trial, 13 patients (56.5%) in rifaximin group and 10 patients (43.5%) in control group. Eradication rate of SIBO was 9/10 (90%) in rifaximin group and 2/6 (33.3%) in control group ($p < 0.05$). In the rifaximin group, gastrointestinal symptom improvement was observed in 4/5 patients aged ≤ 14 years and in 0/5 patients aged > 14 years (p

Authors' conclusions

CF patients show a high prevalence of SIBO, related to a poorer nutritional status. Rifaximin therapy is well tolerated and the results are promising in terms of efficacy in eradicating small intestinal bacterial overgrowth in CF.

<http://dx.doi.org/10.1007/s00535-018-1509-4>

See also

J Gastroenterol. 2018 Sep 19. pii: 10.1007/s00535-018-1509-4. doi: 10.1007/s00535-018-1509-4.

Keywords

Rifaximin; other anti-bacterial agents; Anti-Bacterial Agents; pharmacological_intervention;