

primary studies - published, non RCT

Short-term nutritional supplementation during management of pulmonary exacerbations in cystic fibrosis: a controlled study, including effects of protein turnover.

Code: PM3136639

Year: 1988 **Date:** 1989

Author: Shepherd RW

Participants

1st study: 19 adult CF patients (12 patients completing the study); 2nd study: 10 CF patients

Interventions

1st study: two month programme of home exercise using a cycle ergometer; 2nd study: physiotherapy or exercise

Outcome measures

peak work capacity, maximum oxygen consumption, maximum minute ventilation, sputum weight

Main results

in the 1st study peak work capacity, maximum oxygen consumption, and maximum minute ventilation had increased significantly by the end of the exercise programme; the increase in daily sputum weight (from 24 to 37 g) was not significant ($p = 0.055$). In the second study more sputum was expectorated during and after physiotherapy than during and after exercise (9.8 v 4.0 g).

Authors' conclusions

Exercise may have a role in aiding sputum expectoration in patients with cystic fibrosis but should not be considered as a replacement of physiotherapy.

<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/224/CN-00055224/frame.html>

See also

The American journal of clinical nutrition YR: 1988 VL: 48 NO: 2

Keywords

Adult; exercise; non pharmacological intervention - devices OR physiotherapy; Home; cycle ergometer; Training; Chest physiotherapy;