
primary studies - published RCT

Long-term azithromycin use is not associated with QT prolongation in children with cystic fibrosis.

Code: PM33246911

Year: 2020 **Date:** 2020

Author: Magaret AS

Study design (if review, criteria of inclusion for studies)

Secondary analysis of a placebo-controlled, clinical trial

Participants

Children with CF. 221 study participants with a median of 18 months follow-up

Interventions

Chronic Azithromycin (AZM) taken thrice-weekly for a planned 18 months.

Outcome measures

Safety assessments using electrocardiogram (ECG) occurred at study enrollment, and then after 3 weeks and 18 months of participation.

Main results

Increased corrected QT interval (QTc) of ≥ 30 msec was rare, at 3.4 occurrences per 100 person-years; and incidence of QTc prolongation was no higher in the AZM arm than the placebo arm (1.8 versus 5.4 per 100 person-years). No persons experienced QTc intervals above 500 msec. Long-term chronic AZM use was not associated with increased QT prolongation.

Authors' conclusions

Long-term azithromycin use is not associated with QT prolongation in children with cystic fibrosis.

<http://dx.doi.org/10.1016/j.jcf.2020.11.005>

See also

J Cyst Fibros. 2020 Nov 24:S1569-1993(20)30908-5. doi: 10.1016/j.jcf.2020.11.005.

Keywords

Adolescent; Adult; Anti-Bacterial Agents; Azithromycin; Bacterial Infections; Infection; pharmacological_intervention; Pneumonia; Respiratory Tract Diseases; Respiratory Tract Infections; Macrolides; Anti-Inflammatory Agents; Anti-Inflammatory Agents - excl Steroids;