

primary studies - published RCT

Role of Tris-CaEDTA as an adjuvant with nebulised tobramycin in cystic fibrosis patients with Pseudomonas aeruginosa lung infections: A randomised controlled trial.

Code: PM33341406

Year: 2020 **Date:**

Author: Puvvadi R

Study design (if review, criteria of inclusion for studies)

Randomized controlled trial

Participants

22 adult patients with CF

Interventions

Progressive muscle relaxation (PMR); standard pulmonary rehabilitation (PR). Patients were randomly assigned to either the intervention group (PR + PMR) or the control group (PR only).

Outcome measures

Assessments were performed at baseline, after 21 days of intervention, and at the 48-day follow-up. Outcome measures included the CFQ-R for quality of life, the HADS for mental health, and the PSQI for sleep.

Main results

Compared to the control group, participants who practiced PMR experienced significant reductions in anxiety ($p = 0.05$) and depression ($p = 0.02$) at the final assessment. A significant improvement in sleep quality was also observed (p

Authors' conclusions

Integrating PMR into pulmonary rehabilitation programs may be an effective strategy for improving mental health and sleep in patients with CF. Due to its accessibility and ease of implementation, PMR could be adopted as a complementary method in the holistic care of these patients.

<http://dx.doi.org/10.1016/j.jcf.2020.12.004>

See also

J Cyst Fibros. 2020 Dec 16:S1569-1993(20)30936-X. doi: 10.1016/j.jcf.2020.12.004.

Keywords

non pharmacological intervention - devices OR physiotherapy; non pharmacological intervention - psycho-soc-edu-org; Relaxation Therapy; Behavioural interventions;