

primary studies - published, non RCT

Heterogeneous liver on research ultrasound identifies children with cystic fibrosis at high risk of advanced liver disease.

Code: PM37032248

Year: 2023 **Date:**

Author: Siegel MJ

Study design (if review, criteria of inclusion for studies)

Prospective multicenter case-controlled cohort study.

Participants

Children with pancreatic insufficient cystic fibrosis (CF) aged 3-12 years without known cirrhosis underwent screening US.

Interventions

This study examines whether heterogeneous (HTG) pattern on liver ultrasound (US) identifies children at risk for advanced cystic fibrosis liver disease (aCFLD). Participants with HTG were matched (by age, Pseudomonas infection status and center) 1:2 with participants with normal (NL) US pattern. Clinical status and laboratory data were obtained annually and US bi-annually for 6 years.

Outcome measures

Primary endpoint was development of nodular (NOD) US pattern consistent with aCFLD.

Main results

722 participants underwent screening US, with 65 HTG and 592 NL. Final cohort included 55 HTG and 116 NL with 1 follow-up US. ALT, AST, GGTP, FIB-4, GPR and APRI were higher, and platelets were lower in HTG compared to NL. HTG had a 9.5-fold increased incidence (95% confidence interval [CI]:3.4, 26.7, p

Authors' conclusions

Research US finding of HTG identifies children with CF with a 30-50% risk for aCFLD. A score based on US pattern, age and GPR may refine the identification of individuals at high risk for aCFLD.

<http://dx.doi.org/10.1016/j.jcf.2023.03.019>

See also

J Cyst Fibros. 2023 Apr 7:S1569-1993(23)00083-8. doi: 10.1016/j.jcf.2023.03.019.

Keywords

ultrasound; diagnostic procedures; non pharmacological intervention - diagn;