

primary studies - published RCT

The application of the self-care deficit nursing theory in adolescents with cystic fibrosis: A randomized controlled study.

Code: PM38490107

Year: 2024 Date:

Author: Torun T

Study design (if review, criteria of inclusion for studies)

Randomized controlled study

Participants

30 adolescents with CF

Interventions

Self-Care Deficit Nursing Theory (SCDNT). A total of seven home visits were conducted with the adolescents in the intervention group, during which individualized care plans were applied based on the Self-Care Deficit Nursing Theory.

Outcome measures

The self-care needs of the adolescents in the control group were identified based on data collected during two home visits conducted at the onset of the study and after 4.5 months. RESULTS: As a result of the application of nursing interventions based on the SCDNT, a statistically significant increase was noted in the self-care knowledge and skills of the adolescents in the intervention group (p

Main results

AB - PURPOSE: The present study evaluates the effects of the application of Self-Care Deficit Nursing Theory (SCDNT) on the self-care knowledge and skills in adolescents with Cystic Fibrosis. DESIGN AND METHODS: This randomized controlled study included 30 adolescents who were equally divided into the intervention and control groups. The data were collected by using the Sociodemographic Data Form and the Self-Care Knowledge and Skills Evaluation Forms for Adolescents with Cystic Fibrosis. A total of seven home visits were conducted with the adolescents in the intervention group, during which individualized care plans were applied based on the Self-Care Deficit Nursing Theory. The self-care needs of the adolescents in the control group were identified based on data collected during two home visits conducted at the onset of the study and after 4.5 months. RESULTS: As a result of the application of nursing interventions based on the SCDNT, a statistically significant increase was noted in the self-care knowledge and skills of the adolescents in the intervention group (p

Authors' conclusions

The application of nursing interventions based on the SCDNT was effective in enhancing the self-care knowledge and skills of adolescents with Cystic Fibrosis. PRACTICE IMPLICATIONS: Nurses can benefit from Orem's Self-Care Deficit Nursing Theory in the design and application of the individualized care of adolescents with Cystic Fibrosis to improve their self-care practices.

<http://dx.doi.org/10.1016/j.pedn.2024.03.011>

See also

J Pediatr Nurs. 2024 Jul-Aug;77:96-105. doi: 10.1016/j.pedn.2024.03.011. Epub 2024 Mar 14.

Keywords

non pharmacological intervention - psycho-soc-edu-org;