

primary studies - published RCT

## **A double-blind study of the use of acetylcysteine in patients with cystic fibrosis.**

**Code:** PM5326796

**Year:** 1966 **Date:** 1966

**Author:** Howatt WF

### **Study design (if review, criteria of inclusion for studies)**

Single centre in USA. 4-month trial period, participants receiving interventions for 1 month at a time. No washout period. No dropouts. Data was analysed by ITT, however only limited data was presented in the paper. Randomized, double-blind, controlled, cross-over design.

### **Participants**

8 CF participants (3 male). Age range 6 - 22 years (mean 12.6). Clinical status ranged from "excellent" to "moderate" (based on the method of Shwachman and Kulczycki). 4 participants stopped using nebulized isoproterenol, antibiotics, 3% saline during the trial period, whereas the other 4 participants did not alter their pre-trial therapy. None of the participants had ever received NAC before the trial.

### **Interventions**

Nebulized 20% NAC tid, versus nebulized 2% NAC tid.

### **Outcome measures**

Subjective improvement in sputum thickness and ability to expectorate, PFTs (VC, PEF, PIF, E50, FEV1, SBO), adverse reactions.

<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/587/CN-00000587/frame.html>

### **See also**

University of Michigan Medical Center journal YR: 1966 VL: 32 NO: 2

### **Keywords**

Acetylcysteine; Airway clearance drugs -expectorants- mucolytic- mucociliary-; pharmacological\_intervention; thiols; Respiratory System Agents;