

primary studies - published RCT

Improving the ketchup bottle method with positive expiratory pressure, PEP, in cystic fibrosis.

Code: PM6381081

Year: 1984 **Date:** 1984

Author: Falk M

Study design (if review, criteria of inclusion for studies)

randomised cross-over design

Participants

14 patients with cystic fibrosis

Interventions

Treatment A consisted of postural drainage, percussion and vibration; treatment B of postural drainage and periodic application of a face mask with positive expiratory pressure (PEP); treatment C of PEP in the sitting position; treatment D of the forced expiration technique in the sitting position.

Outcome measures

sputum expectorated. Skin oxygen tension, PSO_2 was monitored continuously during and for 35 min after treatment. patient preference

Main results

In terms of sputum expectorated, treatments B and C were superior to treatment D and especially to treatment A (p less than 0.05). A substantial and prolonged decay in PSO_2 was observed during treatment A, quite different from other patterns seen. During and even following treatment C, an increase in PSO_2 was noted. PEP was well accepted by the patients, who preferred treatment C, and we suggest it is incorporated in chest physical therapy regimens if the therapeutic objective is to increase expectoration.

<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/329/CN-00035329/frame.html>

See also

European journal of respiratory diseases YR: 1984 VL: 65 NO: 6

Keywords

Adolescent; Airway clearance technique; Drainage; Food; non pharmacological intervention - devices OR physiotherapy; non pharmacological intervention - diet; Postural Drainage; Vibration; Positive-Pressure Respiration- PEP- pep mask; forced expiration technique; percussion; Chest physiotherapy; oscillating devices; Active Cycle of Breathing Technique -ACBT-;