

---

*primary studies - published, non RCT*

## **Cough versus chest physiotherapy. A comparison of the acute effects on pulmonary function in patients with cystic fibrosis.**

**Code:** PM6703478

**Year:** 1984 **Date:** 1984

**Author:** de Boeck C

### **Participants**

9 patients with cystic fibrosis who had moderate airways obstruction

### **Interventions**

Vigorous cough and complete chest physiotherapy

### **Outcome measures**

Pulmonary function tests were performed before and 1 h after treatment on consecutive mornings.

### **Main results**

The time spent coughing and in sputum production were equivalent for both treatments. There were no significant differences in functional response to either treatment. Static lung volumes and flows at high lung volumes were not affected. The increases in maximal expiratory flows at 50 and 25% of the vital capacity were 14 +/- 10 SEM% and 22 +/- 11 SEM%, respectively, after cough and 18 +/- 8% and 25 +/- 13% after chest physiotherapy. These responses did not correlate with the volume of sputum produced.

### **Authors' conclusions**

Because there was no clear-cut benefit of chest physiotherapy over cough alone, and the latter is easier to perform, cough is an attractive alternate method of treatment.

<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/131/CN-00208131/frame.html>

### **See also**

American Review of Respiratory Disease YR: 1984 VL: 129 DE: RCT NO: 1

### **Keywords**

Adolescent; Child; non pharmacological intervention - diagn; Respiratory Tract Diseases;