

primary studies - published RCT

Comparison of a high lipase pancreatic enzyme extract with a regular pancreatin preparation in adult cystic fibrosis patients.

Code: PM7696449

Year: 1994 Date: 1998

Author: Gan KH

Study design (if review, criteria of inclusion for studies)

RCT

Participants

16 patients (eight girls and eight boys) with cystic fibrosis, 4 to 20 years of age (12 ± 1.3 , mean \pm SEM), who were pancreatic insufficient

Interventions

two 6-day regimens of nocturnal continuous enteral feedings offered in random order. Forty-four percent of the total daily energy was consumed as enteral tube feedings, and the remaining oral dietary intake remained constant throughout the study.

Outcome measures

Seventy-two-hour fecal collections from each study period were analyzed for total fat, long-chain fatty acids (LCFAs), medium-chain fatty acids (MCFAs), and nitrogen.

Main results

Fat absorption was $80.2\% \pm 2.9\%$ and $82.3\% \pm 3.1\%$ ($p = 0.58$) for the semi-elemental and nonelemental formulas, respectively. Similarly, the coefficient of absorption of LCFAs was $69.5\% \pm 4.5\%$ and $79.6\% \pm 3.4\%$ ($p = 0.30$) for the semi-elemental and nonelemental formulas. Malabsorption of MCFAs contributed minimally to total fat malabsorption. There was no difference between formulas for MCFA or for nitrogen absorption, $83.7\% \pm 1.9\%$ and $87.4\% \pm 1.4\%$, $p = 0.48$. All patients tolerated all feedings, and weight gained was 1.8 ± 0.3 kg with no difference in weight gain between the trials.

Authors' conclusions

A nonelemental formula, with enzyme replacement, is absorbed as well as a predigested formula in patients with CF who are pancreatic insufficient.

<http://dx.doi.org/10.1111/j.1365-2036.1994.tb00337.x>

See also

Aliment Pharmacol Ther. 1994 Dec;8(6):603-7.

Keywords

Adolescent; Adult; Child; Enteral Nutrition; Food; non pharmacological intervention - diet; Supplementation; Malabsorption; Nutrition Disorders; Oral;