

---

*primary studies - published, non RCT*

## **Conventional in-vitro fertilization versus intracytoplasmic sperm injection for patients requiring microsurgical sperm aspiration.**

**Code:** PM7836522

**Year:** 1994 **Date:** 1994

**Author:** Silber SJ

### **Participants**

men with CBAVD or irreparable obstructive azoospermia

### **Interventions**

ICSI vs IVF

### **Outcome measures**

fertilization, % normal embryos, pregnancy rate

### **Main results**

ICSI (using epididymal or testicular spermatozoa) from men with CBAVD or irreparable obstructive azoospermia, achieved good fertilization and normal embryos in 82% of cases, compared to 19% with conventional IVF. There was an overall fertilization rate of 45%, with 85% progressing to normally cleaving embryos using ICSI, compared to 6.9% using conventional IVF. The pregnancy rate with ICSI/MESA was 47% per stimulated cycle (normal delivery rate was 30%), compared to 4.5% with conventional IVF. These results were achieved in patients who had consistently failed to fertilize in previous cycles with MESA and conventional IVF.

<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/683/CN-00109683/frame.html>

### **See also**

Human reproduction (Oxford, England) YR: 1994 VL: 9 NO: 9

### **Keywords**

Adult; fertilization; Gonadal Disorders; Infertility- Male; Microinjections; non pharmacological intervention - genetic& reprod; Pregnancy; surgery; non pharmacological intervention - surg;