

primary studies - published RCT

## Eicosapentaenoic acid in cystic fibrosis: evidence of a pathogenetic role for leukotriene B4.

**Code:** PM8102430

**Year:** 1993 **Date:** 1993

**Author:** Lawrence R

### Study design (if review, criteria of inclusion for studies)

6-week cross-over study planned.

### Participants

19 adolescents and adults diagnosed with CF on genotype, sweat test, or clinically (median age 17 years, age range 12 years to 26 years), 11 male, 5 female. 3 participants excluded due to requiring a course of corticosteroids for asthma attacks. All were recruited within 4 weeks of hospitalisation for acute respiratory infection and judged to be in optimum condition for disease stage. All were *Pseudomonas aeruginosa* colonised and produced at least 5 ml sputum daily. Initial randomisation gave groups with comparable baseline characteristics except for age. The treatment group also had significantly greater weight, peripheral blood leucocyte and neutrophil counts.

### Interventions

2.7 g EPA daily compared with identical olive oil placebo capsules, over 6 weeks.

### Outcome measures

Outcomes included in this review: number of people experiencing adverse events; number of deaths; changes in haematological and growth indices; changes in lung function; changes in in-vitro neutrophil chemotaxis.

### Main results

EPA was well tolerated and resulted in a significant reduction in sputum volume (median change with EPA -10 mL/day, placebo 0;  $p = 0.015$ ), and improvements in Schwachman score (EPA 5%, placebo 0;  $p = 0.034$ ), forced expiratory volume in 1 s (EPA 0.25 L, placebo -0.1 L;  $p = 0.006$ ), and vital capacity (EPA 0.6 L, placebo 0;  $p = 0.011$ ). Relative chemotaxis of circulating neutrophils to LTB<sub>4</sub> increased from a subnormal baseline of 4 (median; range 0-10) microns/30 min before treatment, to a near normal value of 11 (5-18) microns/30 min after EPA. Relative chemotaxis to LTB<sub>4</sub> of patients taking placebo did not change: the difference in response was highly significant ( $p = 0.001$ ). Specific reduction of neutrophil chemotaxis to LTB<sub>4</sub> is a sensitive assay of chronic in-vivo exposure to LTB<sub>4</sub>.

### Authors' conclusions

LTB<sub>4</sub> has a pathogenetic role in the lung damage of cystic fibrosis. Longer-term clinical trials of EPA are warranted in a larger number of cystic fibrosis patients.

[http://dx.doi.org/10.1016/0140-6736\(93\)91594-C](http://dx.doi.org/10.1016/0140-6736(93)91594-C)

### See also

Lancet YR: 1993 VL: 342 NO: 8869

### Keywords

Adolescent; Adult; Bacterial Infections; Child; Infection; non pharmacological intervention - diet; pharmacological\_intervention; placebo; *Pseudomonas aeruginosa*; *Pseudomonas*; Respiratory Tract Diseases; Respiratory Tract Infections; Supplementation; essential fatty acids; omega-3; Eicosapentaenoic acid -EPA-; Capsules;