

primary studies - published RCT

## A comparison of bronchial drainage treatments in cystic fibrosis.

Code: PM8905888

Year: 1996 Date: 2000

Author: Klufft J

### Study design (if review, criteria of inclusion for studies)

randomized trial

### Participants

60 patients with cystic fibrosis, chronically infected with *P aeruginosa*

### Interventions

two treatment arms (elective or symptomatic) and yearly follow-up

### Outcome measures

The major end points were changes in forced expiratory volume in one second (FEV(1)) and forced vital capacity (FVC). Survival was a secondary end point.

### Main results

Patients in the symptomatic group received a mean of three antibiotic treatments each year and those in the elective group received four antibiotic treatments during each year of the study. No significant differences in FEV(1) and FVC were found between the two groups after three years. There was a statistically non-significant higher rate of deaths in the elective group (n = 4), three of which were associated with *B cepacia* infection, compared with the symptomatic group (n = 0).

### Authors' conclusions

This study did not demonstrate an advantage of a policy of elective antibiotic treatment over symptomatic treatment in patients with cystic fibrosis chronically infected with *Pseudomonas* species.

[3.0.CO;2-P" target=" blank">http://dx.doi.org/10.1002/\(SICI\)1099-0496\(199610\)22:4<271::AID-PPUL7>3.0.CO;2-P](http://dx.doi.org/10.1002/(SICI)1099-0496(199610)22:4<271::AID-PPUL7>3.0.CO;2-P)

### See also

Pediatr Pulmonol. 1996 Oct;22(4):271-4.

### Keywords

Anti-Bacterial Agents; Bacterial Infections; Infection; pharmacological\_intervention; *Pseudomonas aeruginosa*; *Pseudomonas*; Respiratory Tract Diseases; Respiratory Tract Infections;