

primary studies - published RCT

Intravenous colistin sulphomethate in acute respiratory exacerbations in adult patients with cystic fibrosis.

Code: PM9487348

Year: 1997 **Date:** 2001

Author: Conway SP

Study design (if review, criteria of inclusion for studies)

double-blind placebo-controlled crossover design.

Participants

7 prepubertal CF children aged 9.6 to 13 years (5 boys and 2 girls) .

Interventions

CF patients were treated with placebo or IGF-1 for 6 months. After a 6-month washout period, patients received the alternative therapy for 6 months

Outcome measures

The primary outcome measure was linear growth rate. Secondary outcome measures were changes in body mass index, body composition determined by dual energy x-ray absorptiometry, forced expiratory volume (FEV(1)), and the blood glucose/insulin ratio.

Main results

The mean height z score at baseline was -1.5 ± 0.8 . At entry, the mean serum IGF-1 level was 124 ± 25 ng/mL (normal range, 110-771 ng/mL). With treatment, mean serum IGF-1 levels increased twofold to threefold for all patients. The half-life for IGF-1 was 10.3 hours. We observed no significant difference in linear growth rate, weight gain, rate of accretion of lean body mass, or mean FEV(1) during treatment with IGF-1 compared with placebo. The glucose/insulin ratio, an indirect index of insulin sensitivity, was significantly increased with IGF-1 treatment compared with placebo (P

Authors' conclusions

Treatment with IGF-1 for 6 months did not promote linear growth in prepubertal children with CF. However, the glucose/insulin ratio was increased without changing blood glucose levels with IGF-1 treatment suggesting increased insulin sensitivity.

<http://dx.doi.org/10.1136/thx.52.11.987>

See also

Thorax. 1997 Nov;52(11):987-93.

Keywords

Adolescent; Child; Insulin-Like Growth Factor Binding Protein 3; Insulin-Like Growth Factor I; Nutrition Disorders; pharmacological_intervention; Malnutrition; Hormones; Growth Hormone;