

primary studies - published, non RCT

Comparison of neoral and sandimmun cyclosporines for de novo lung transplantation in cystic fibrosis patients.

Code: PM9636614

Year: 1998 **Date:** 2020

Author: Mikhail G

Study design (if review, criteria of inclusion for studies)

Qualitative interview study undertaken concurrently with a pilot randomized controlled trial.

Participants

Adults with cystic fibrosis. 14 patients in the intervention arm of the trial, and five members of the multidisciplinary teams offering wider care to patients.

Interventions

A new complex intervention to help adults with cystic fibrosis increase adherence to nebulizer treatments was tested. A qualitative interview study undertaken concurrently with a pilot randomized controlled trial. 25 semi-structured interviews were conducted with three interventionists at two time points,

Outcome measures

behavior change

Main results

The intervention was informed by a theoretical framework of behavior change. There was evidence of the expected behavior change mechanisms of action. There was also evidence of Vassilev's mechanisms of action associated with effective telehealth interventions for self-management support: relationships, visibility, and fit. Patients described how building a relationship with the interventionist through face-to-face visits with someone who cared about them and their progress helped them to consider ways of increasing adherence to medication. Rather than seeing the visibility of adherence data to clinicians as problematic, patients found this motivating, particularly if they received praise about progress made. The intervention was tailored to individuals but there were challenges in how the intervention fitted into some patients' busy lives when delivered through a desktop computer.

Authors' conclusions

Mechanisms associated with effective telehealth interventions for self-management operated within this new intervention. The intervention was modified to strengthen mechanisms of action based on these findings, for example delivery through an app accessed via mobile phones, and then tested in a randomized controlled trial in 19 UK cystic fibrosis centers.

<http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/012/CN-00688012/frame.html>

See also

Transplantation proceedings YR: 1998 VL: 30 NO: 4

Keywords

non pharmacological intervention - diet; non pharmacological intervention - psycho-soc-edu-org; Nutrition Disorders; Behavioural interventions; training; Inhalation OR nebulised;