

primary studies - published RCT

## **Holding the baby: head downwards positioning for physiotherapy does not cause gastro-oesophageal reflux.**

**Code:** PM9817174

**Year:** 1998 **Date:** 1998

**Author:** Phillips GE

### **Study design (if review, criteria of inclusion for studies)**

RCT, cross-over design

### **Participants**

21 infants (age range 1-27 months) with respiratory disorders (CF=11) undergoing lower oesophageal pH monitoring were recruited.

### **Interventions**

Subjects received two physiotherapy episodes in random order, A/B or B/A, 12 h apart. A began the gravity-assisted positioning head downward tip for: right lower lobe, middle lobe, left lower lobe and lingula; then supine with no tip for anterior segments of the upper lobes followed by apical segments of upper lobes in a sitting position. B was in the reverse order. Intermittent chest clapping was carried out for 4 min in each position by a physiotherapist blinded to the pH data.

### **Outcome measures**

lower oesophageal Ph

### **Main results**

pH from baseline was -0.32 (range -2.07 to +1.0) in non-CF subjects (NS) and -0.52 (range -2.7 to +0.52) in CF subjects (p

### **Authors' conclusions**

the head-downward tipped positioning for physiotherapy treatment neither induces nor aggravates gastro-oesophageal reflux. There is no justification for routinely changing the way in which infant physiotherapy is carried out.

<http://dx.doi.org/10.1183/09031936.98.12040954>

### **See also**

Eur Respir J. 1998 Oct;12(4):954-7.

### **Keywords**

Gastrointestinal Diseases; Infant; Newborn; non pharmacological intervention - devices OR physiotherapy; Postural Drainage; screening; Intermittent; clapping; Airway clearance technique; Chest physiotherapy;